

Erratum:

Safety and efficacy of sedation/analgesia administered by the urologist for minimally invasive transurethral procedures

A. Beri, N.J. Mabweesh, M. Sofer, H. Matzkin, J. Chen, A. Greenstein

Ambulatory Surgery 2017;23.3:74–7

An error occurred in the transcription of Table 2, where the ASA grade of patients was incorrectly inserted at the proofing stage of the edition.

The correct table is shown below:

Table 2 Procedure-related measures (n = 77).

Measure	Patients, n
ASA grading	
I	31
II	37
III	9
Self-report pain score (0–10) (n=71)	
1	4
2	1
4	1
Highest blood pressure (mmHg)	
Mean (SD)	152.4 (20.6)
Median (IQR)	151 (140–168)
Lowest blood pressure (mmHg)	
Mean (SD)	123.5 (18.5)
Median (IQR)	124 (110–134)
Highest pulse (bpm)	
Mean (SD)	82.7 (15)
Median (IQR)	81 (72–91)
Lowest pulse (bpm)	
Mean (SD)	67.7 (10.1)
Median (IQR)	65 (60–74)
Lowest oxygen saturation (%)	
Mean (SD)	94.2 (2)
Median (IQR)	94 (93–96)

SD = standard deviation; IQR = interquartile range.

Mark Skues
Editor-in-Chief